



STK 573
Metode Grafik untuk Analisis dan Penyajian Data

Pertemuan 6
Sajian Peubah Kontinu Tunggal

Tim Dosen:
Prof. Dr. Khairil Anwar Notodiputro
Dr. Ir. Aji Hamim Wigena
Dr. Agus M Soleh

Introduction

- exploring distributions of a single continuous variable
- continuous variables as plots
- discrete (categorical) variables as charts

Introduction


- The Dotplot
- The Stemplot
- The Boxplot
- The EDF Plot

The Dotplot





- one-dimensional scatterplots
- referred to as a stripchart
- Original:
 - Replicated observations occur as a single dot
 - If observations occur sufficiently close together the separation between dots is not apparent at a given scale and they can also appear to be a single dot
- function: `stripchart()`

Mass (g)

5.9	32.0	40.0	51.5	70.0	100.0	78.0	80.0
85.0	85.0	110.0	115.0	125.0	130.0	120.0	120.0
130.0	135.0	110.0	130.0	150.0	145.0	150.0	170.0
225.0	145.0	188.0	180.0	197.0	218.0	300.0	260.0
265.0	250.0	250.0	300.0	320.0	514.0	556.0	840.0
685.0	700.0	700.0	690.0	900.0	650.0	820.0	850.0
900.0	1,015.0	820.0	1,100.0	1,000.0	1,100.0	1,000.0	1,000.0



- Variations on the Dotplot:
 - stacked dotplot -> identical observations can be ameliorated
 - open circles -> eliminate smear over a range of values
 - Jittering
 - barcode plot -> dot icon replaced by a vertical line segment



The Stemplot

- or the stem-and-leaf plot
- depicting distributions of a numerical variable
- Features:
 - For data sets of reasonable size, each observed value of the quantitative variable can be listed thereby giving a complete picture of all the individual data points as well as a picture of the distribution

- Features (continued):
 - The stem can be prepared quickly and easily by hand for small data sets
 - It is useful for hand calculation of quantiles and thus is useful in preparing another well-known EDA plot—the boxplot and even a histogram
- Function : `stem()`

The decimal point is 2 digit(s) to the right of the |

0		134578899
1		011222333345555789
2		0235567
3		002
4		
5		16
6		599
7		00
8		2245
9		00
10		0002
11		00

1 | 2: represents 120
leaf unit: 10
n: 56

9	0		034577888
28	1		0111222333344557889
(6)	2		125566
22	3		002
	4		
19	5		15
17	6		589
14	7		00
12	8		2245
8	9		00
6	10		0001
2	11		00

The Boxplot



- box-and-whisker plot
- box-whisker plot
- the boxplot require the concept of a quantile
- Function : `boxplot()`

The plotting convention for the quantile boxplot consists of:

- a rectangular box with edges determined by the lower and upper quartiles;
- the median denoted as a line segment splitting the rectangular box into two adjoining boxes;
- a whisker (that is, a line segment) from Q1 to the minimum; and
- a whisker from Q3 to the maximum.

The plotting convention for the outlier boxplot consists of:

- a rectangular box with edges determined by the lower and upper quartiles;
- the median denoted as a line segment splitting the rectangular box into two adjoining boxes;
- whiskers ending at the adjacent values; and
- open circles to represent points lying beyond the inner fences.





The EDF Plot

- A discrete approximation to the distribution of the random variable X is given by the empirical distribution function (EDF):

$$S_n(x) = \begin{cases} 0 & \text{if } x < x_{(1)}, \\ \frac{i}{n} & \text{if } x_{(i)} \leq x < x_{(i+1)}, \\ 1 & \text{if } x \geq x_{(n)}. \end{cases}$$

- the empirical distribution function is an approximation to the cumulative distribution function of the random variable X so it is also known as the empirical cumulative distribution function (ecdf)

- plotted along the horizontal axis rather than the vertical axis
- their plot is actually the inverse of the empirical cumulative distribution function'
- Function : `ecdf()`, `plot.stepfun()`





Selesai